Funding Science • Finding Solutions • Fuel the Change

Why Guess? Fill Knowledge Gaps With New Insights from Field-based Research

August 1, 2019 10:00am Pacific Standard Time



Webinar Instructions

Attendees will be on mute during the webinar

Please send questions through chat box

Connection questions may be directed to: info@centerforproducesafety.org



Speakers

Dave Corsi

VP of Produce and Floral at Wegmans Food Markets Chair, Center for Produce Safety Board of Directors

Dr. Channah Rock,

Professor and Water Quality Extension Specialist University of Arizona

Dr. Trevor Suslow,

Emeritus Extension Faculty, UC Davis
Vice President, Produce Safety, Produce Marketing Association

Bonnie Fernandez-Fenaroli

Executive Director, Center for Produce Safety



Welcome



VP of Produce and Floral Wegmans Food Markets &

Chair, Center for Produce Safety

Board of Directors



Center for Produce Safety

Mission:

Fund the science, find solutions and fuel the change

Purpose:

Providing the science to support produce safety



Rapid Response Research

- Research opportunities that will be lost if not implemented quickly
- Industry partner require
- Maximum Project Cost \$50,000 maybe larger if cash match is available
- CPS funds available Will provide up to 50% of total project cost as a match to other cash pledges.
- Proposal and Budget must be submitted at the same time.
- Once the proposal and budget is received. A subcommittee of the CPS Technical Committee is formed. A review will be provided within 3-5 days of the proposal receipt.
- The receiving university must be ready to implement a contract quickly.





Dr. Channah Rock Professor and Water Quality Extension Specialist University of Arizona



CPS Rapid Response – Yuma Valley Objectives

 To capitalize on the unique opportunity to study romaine production under real world conditions within close proximity to potentially implicated fields from the 2018 outbreak

 To generate new knowledge that is useful to industry in order to bridge the gap created by the recent outbreak



Scope Of Work

1. Water treatment

2. Persistence of microbial contamination on Romaine after **freeze injury**

3. Industry sampling protocols and pathogen transport through commercial scale harvest



Industry Support

- Two 3 acre plots
- Three varieties of Romaine
 - Del Sol, Sunbelt, 8403
- Furrow and Sprinkler Irrigation*
- Commercial Scale Harvester/Crew
- Staff time/resources



*same canal water source



1. Water Treatment

- Peroxyacetic Acid (PAA)
 - —2 to 6 ppm target concentration*

- 12% Sodium Hypochlorite (NaOCl)
 - —3 to 5 ppm target concentration*



Water Treatment

- Triplicate trials of each sanitizer
- Positive control field plots (3)
- Spiking Strain
 - -TVS353 (ECC ChromAgar + 80ug/ml rif)
 - Isolated from Irrigation Water
 - −10² to 10⁴ CFU Target Concentration






Water Treatment







Water Treatment

- Sample Types
 - Irrigation Water
 - Grab Samples (birds), mMS filters
 - Calculate log reductions
 - Plant Tissue
 - Industry protocols, 'S' and 'Z' (n=60+)
 - Presence/Absence
 - Soil
 - Grab/Composite samples
 - Presence/Absence






Irrigation Water

Concentration (CFU)	492
Detection Limit	1
Log Reduction	2.69

- 1st and last sprinkler head
 - -Log reduction >2.69 for both PAA and NaOCI*



Enrichment



Plant Tissue

PAA


- "S" pattern = 50% positive
- "Z" pattern = 25% positive

Sodium Hypochlorite

- "S" pattern = 0% positive
- "Z" pattern = 0% positive

Positive Control

- "S" pattern = 100%
- "Z" pattern = 96%








Soil

- PAA
 - -Zones 1-5 (66% positive)

- Sodium Hypochlorite
 - -Zones 1-5 (33% positive)

- Positive Control
 - -Zones 1-5 (66% positive)






2. Freeze Damage

"What is the impact of freeze damage on harborage of bacteria?"




Freeze Damage

■ Damaged ■ Un-damaged



Freeze Damage




3. Field Contamination/Harvest

Three "Contamination" Event Types


- 1. Contamination "Pulse" Furrow Irrigation
- 2. Animal Intrusion
- 3. Aerial Application



Contamination "Pulse" Furrow Irrigation











Animal Intrusion







Aerial Application



 Mimic pesticide application

Known/measured application rates

Highly controlled



3. Field Contamination/Harvest

- Sample type
 - —Raw Product pre-harvest (field)
 - Workers gloves (swabs)
 - —Harvest tools (swabs)
 - —Table surface (swabs)
 - –Conveyer belt (swabs)
 - —Elevator (swabs)
 - Raw Product post-harvest (disposable bins)



Commercial Scale Harvester

40+ person crew


9 harvesting "stations"



Commercial Scale Harvester










Commercial Scale Harvester













Raw Product Pre-Harvest

Animal

Furrow Irrigation

ation Intrusion




Bin 4

Bin 3

Bin 2

Bin 1



Furrow Irrigation

	Stop 1	Stop 2	Stop 3	Stop 4	Stop 5
Gloves	0%	20%	0%	0%	0%
Knife	0%	20%	20%	0%	0%
Table	0%	20%	0%	0%	0%
Belt	0%	0%	0%	0%	0%
Elevator	0%	0%	0%	0%	0%



Animal Intrusion

	Stop 1	Stop 2	Stop 3	Stop 4	Stop 5
Gloves	0%	40%	60%	40%	40%
Knife	0%	40%	60%	0%	20%
Table	0%	60%	80%	60%	60%
Belt	0%	20%	0%	0%	0%
Elevator	0%	100%	100%	100%	100%



Aerial Application

	Stop 1	Stop 2	Stop 3	Stop 4	Stop 5
Gloves	0%	80%	80%	80%	100%
Knife	0%	80%	80%	80%	80%
Table	0%	80%	80%	80%	80%
Belt	0%	100%	80%	80%	0%
Elevator	0%	100%	100%	100%	100%



Raw Product Post-Harvest Bins

Furrow Irrigation Animal Intrusion <u>Aerial</u> Application



What we learned...



 Sanitizers were effective at reducing microbial loading in irrigation water

 Bacteria may survive water treatment and persist in soil and on plant surface

Sample volume matters!



What we learned...

 At high concentrations, bacteria were able survive on "freeze-damaged" romaine



Variety type had minimal impact

 At lower concentrations, "freeze damaged" romaine may have the ability to harbor bacteria extended periods of time



What we learned...

 Pre-harvest sampling protocols on 1 acre plots detected contamination, but post harvest sampling was more consistent

Contamination by furrow irrigation was detectable through harvest

Harvest practices aid in contamination transfer

 Animal Intrusion and Aerial Application contamination events result in pervasive contamination

Getting the work done!

- 50+ UA Students and Staff
- Industry Support!!
- Initiated field work Feb. 18th
- 3 straight weeks in field
- Complete lab work (culture)
 April 8th
- Molecular work on-going
- 2000+ Samples































Dr. Trevor Suslow Emeritus Extension Faculty,
UC Davis

VP Produce Safety Produce Marketing Association





Yuma Dust Storm Microbial Community



ORIGINAL ARTICLE

Leaf microbiota in an agroecosystem: spatiotemporal variation in bacterial community composition on field-grown lettuce

Gurdeep Rastogi¹, Adrian Sbodio², Jan J Tech¹, Trevor V Suslow², Gitta L Coaker¹ and Johan HJ Leveau¹

¹Department of Plant Pathology, University of California, Davis, CA, USA and ²Department of Plant Sciences, University of California, Davis, CA, USA

Yuma typical Romaine microbial community

WHY GUESS? – dust storms change the microbial community and bring new 'transients' to crops







carrier down towards basal sector

Organic dust accumulates at basal leaf whorl





Recovery of *E. coli_{rif}* in Feclone from Romaine Lettuce Plants with Overhead Sprinkler Irrigation

Air Temperature During Field Trial

Hour	High (°F)	Low (°F)
0	86	52
24	80	62
48	87	68
72	93	65
96	102	68

Analysis Sectioning of Romaine Lettuce





At 96h, after three irrigation events, there is a concentration of generic *E. coli* at the bottom of the Romaine Lettuce plant at sections C and D.



5 People Died From Eating Lettuce, But Trump's FDA Still Won't Make Farms Test Water For Bacteria

09/27/2018 11:26 am ET | Updated Sep 27, 2018



PAUL SAKUMA/ASSOCIATED PRESS

A worker harvests romaine lettuce in Salinas, Calif., in August 2007. After an E. coli outbreak on spinach killed three people in 2006, California and Arizona growers of leafy greens voluntarily agreed to test their irrigation water and take other safety steps.

By Elizabeth Shogren and Susie Neilson



Furrow Irrigation: Majority of implicated Yuma fields furrow irrigated to maturity

Potential for irrigation water contact across the planted bed and onto lower leaves and surrounding soil











Project Outline

- Furrow water with generic *E. coli_{Rif}* sachets
 - Trial 1 Low Inoculation Log 1
 - Trial 2 High Inoculation Log 5
- Bed-edge cross-contact with lowest leaves
- Furrow dry to walkable (~ 3 days)
- Harvest romaine lettuce heads
- Test knife and platform
- Sample non-harvest leaves
- Sample head (market head) to platform
- Cut and invert for 20 minutes before removing from field (Inverted Leaves)
- Sample furrow soil.



E. coli Persistence in Furrow Soil

Low Log 1 Inoculation

E. coli recovery from furrow soil samples in Trial 1 at 8 days

Positive for
E.coli- rif

Key

Negative for E.coli - rif

Furrow Soil Sample	Enrichment
1	+
2	+
3	+
4	+
5	+
6	+
7	+
8	+
9	+
10	+

High Log 5 Inoculation

E. coli recovery from furrow soil samples in Trial 2 at 8 days

Furrow Soil Sample	Enrichment
1	+
2	+
3	+
4	+
5	+
6	+
7	+
8	+
9	+
10	+



E. coli Recovery from Romaine Lettuce Lowest Non-market Leaves

Low Log 1 Inoculation

E.coli recovery from leaves that came in contact with furrow water for Trial 1

Lowest Leaves	Reps	LogCFU/g	Average LogCFU/g	SD	Enrichment
	1	0.86			+
Bed 1	2	0.08		0.48	+
	3	0.95			-
	1	0.26			+
Bed 2	2 0.89	0.20	0.58	+	
beu z	3	Below	0.29	0.29	
	5	LOD			-
	1	-0.22			+
Bed 3	2	0.26	0.14	0.32	+
	3	0.38			+

High Log 5 Inoculation

E.coli recovery from leaves that came in contact with furrow water for Trial 2

Lowest Leaves	Reps	LogCFU/g	Average Log CFU/g	SD
	1	4.43		
Bed 1	2	4.49	4.33	0.22
	3	4.09		
	1	4.00		
Bed 2	2	3.35	3.65	0.33
	3	3.60		
	1	3.50		
Bed 3	2	2.78	2.89	0.57
	3	2.38		

Limit of Detection (LOD) = -Log 0.27/g



E. Coli Recovery from Market-Head Romaine Lettuce Leaves (not inverted)

Low Log 1 Inoculation

E. coli recovery of marketable Romaine Leaves in Trial 1

Market Leaves	Reps	LogCFU/g	Enrichment
	1	Below LOD	-
Bed 1	2	Below LOD	-
	3	Below LOD	-
	1	Below LOD	-
Bed 2	2	Below LOD	-
	3	Below LOD	-
Bed 3	1	Below LOD	-
	2	Below LOD	-
	3	Below LOD	-

Limit of Detection (LOD) = -Log0.27/g

High Log 5 Inoculation

E. coli recovery of marketable Romaine Leaves Trial 2

Market Leaves	Reps	LogCFU/g	Enrichment
	1	Below LOD	-
Bed 1	2	Below LOD	-
	3	Below LOD	-
	1	Below LOD	+
Bed 2	2	Below LOD	+
	3	Below LOD	+
	1	-0.22	
Bed 3	2	Below LOD	+
	3	0.38	



E. coli Recovery from Inverted Market-Head Romaine Lettuce Leaves

High Log 5 Inoculation

Inverted Romaine	Reps	Log CFU/g	Average Log CFU/g	SD
	1	1.56		
Bed 1	2	1.44	1.43	0.13
	3	1.30		
	1	2.42		
Bed 2	2	2.31	2.36	0.06
	3	2.36		
	1	2.68		
Bed 3	2	2.57	2.66	0.08
	3	2.73		
Bed 3	1	2.36		
	2	2.34	2.33	0.02
	3	2.31		

E. coli was recovered from top leaves of heads inverted onto the basal leaves for 20 minutes.

Cross Contamination - Knife

Low Log 1 Inoculation

E. coli recovery from swabs of knife used during harvesting in Trial 1

Lettuce Harvested	Knife
16	_
32	-
48	-
64	+
80	+
96	+
112	-
128	+
144	-
160	+
176	+
192	-
208	-
224	-
240	+

High Log 5 Inoculation

E. coli recovery from swabs of knife used during harvesting in Trial 2

Lettuce Harvested	Knife
12	+
24	+
36	+
48	+
60	+
72	+
84	+
96	+
108	+





Cross Contamination - Surfaces

Low Log 1 Inoculation

E. coli recovery from swabs of surfaces that came in contact with romaine lettuce heads (metal tray and cutting board) in

Trial 1

Lettuce Harvested	Metal Tray	Cutting Board
8	-	-
16	-	-
24	-	+
32	-	-
40	-	-
48	-	-
56	-	-
64	-	-
72	-	-
80	-	-
88	-	1
96	_	_
104	+	-
112	-	_
120	-	+

High Log 5 Inoculation

E. coli recovery from swabs of surfaces that came in contact with romaine lettuce heads (metal tray and cutting board) in

Trial 2

Lettuce Harvested	Metal Tray	Cutting Board
6	-	+
12	+	+
18	+	+
24	+	+
30	+	+
36	+	+
42	+	+
48	+	+
54	+	+



What are the integrated solutions?

- Deadhead buffer at field ends
- Raise bed height
- Modify cut-to-ground
- Change pre-harvest sampling protocols
 - Testing is not a food safety plan, but...
- Your input would be to ...





Bonnie Fernandez-Fenaroli Executive Director Center for Produce Safety



www.centerforproducesafety.org

CPS CENTER for PRODUCE SAFETY

HOME | CONTACT US

ABOUT


GRANT OPPORTUNITIES FUNDED RESEARCH PROJECTS

NEWS -

SUPPORT

ANNUAL RESEARCH SYMPOSIUM

RESOURCES



Change begins with a vision

2018 CPS Annual Report

Knowledge Transfer Task For

Spreading CPS research knowledge to all

RFP ANNOUNCEMENT!

Water Treatment Baseline Program



Funded Research Projects





Questions? Please use chat box





THANK YOU.

YOUR TIME AND INTEREST IS APPRECIATED

SEND ADDITIONAL QUESTIONS TO:

INFO@CENTERFORPRODUCESAFETY.ORG

